

Jeremy W. Denning, M.D.

Scoliosis/Lumbar Fusion: Post-Operative Instructions

Incision Care

- If your incision is closed with dermabond, there is no need to cover it. If your incision is closed with staples, your dressing will be a gauze pad and tape. It should be changed daily while staples are in place.
- You may shower, but no tub baths. Cover the incision to keep dry while staples are in place.

Pain

You may still have some pain, numbness or tingling in your back, hips or legs. This should decrease gradually as the nerves heal. An ice pack may be used on the incision to reduce swelling and pain.

Brace Instructions

- You must wear your brace at all times when out of bed.
- You must take it off to shower.
- Your physician will determine when you are ready to wean out of your brace, but you are usually wearing the brace for 8 to 12 weeks.

Call Our Office

If you...

- Feel severe pain, weakness or numbness in your leg(s).
- Notice excessive swelling or increased redness around your incision.
- If bloody drainage requires 2 or more bandage changes in a single day.
- If any clear fluid drainage comes from wound.
- Have a fever of 101 degrees or greater.
- Have a new problem controlling your bladder or bowel.

Medications

■ You will be given a prescription for pain medicine when you are discharged from the hospital.

Take as directed on the prescription. **Do not drink alcoholic beverages while taking this medication.**

■ You will be given a prescription for muscle relaxants (to reduce spasms or tightness in your back).

This medication may make you drowsy. **Do not drink alcoholic beverages while on this medication.**

■ Medications will be refilled during normal business hours (8:30 am – 5:00 pm, M-Thurs, 9:30 am – noon Fri), as the physician on call cannot renew prescriptions.

Jeremy W. Denning, M.D.

Scoliosis/Lumbar Fusion: Post-Operative Instructions(cont.)

Activity

First Week:

- Avoid bending or twisting from the waist.
- Do not lift over 5 pounds until released to do so.
- Limit auto travel. Passenger only. No driving.
- Walk around the house for first couple of days after surgery. Then you may begin to walk outside for short periods 1 to 2 blocks per day. Do not overdo it.
- Avoid house and yard work. You may cook but do not lift more than 5 pounds until released to do so.
- You may shower, but no tub baths.
- To sleep you may lie on your back or side, but not on your abdomen.
- Take a nap or at least rest in the morning and again in the afternoon.

Second Week:

- Increase walking distance as tolerated, working up to 1 mile per day.
- Take a nap or at least rest in the morning and again in the afternoon.
- Continue to avoid house and yard work.
- You may begin driving short distances, no more than 30 minutes. Make certain your seat, mirrors and restraints are properly adjusted.
- Continue bending, twisting and lifting instructions.

Third Week:

- Increase walking distance as tolerated.
- You may be up and about as tolerated.
- Begin light housework. Do not sweep or vacuum. Stop for frequent breaks. Use good body mechanics.
- Continue bending, twisting, and lifting instructions.

Follow-Up Appointment

Your first post-op appointment should be 2-3 weeks from your surgery. Please call early to get a convenient appointment time. Your next appointment will be 6-8 weeks later.

Return to Work

Return to work varies with the type of occupation and will be discussed at your follow-up appointment.